

Effect of Thyme Essential Oil On Drug-Resistant *Pseudomonas* Isolated From Cystic Fibrosis Patients

Ghazal M A Bishal

Department of Chemistry, Faculty of Science, Arak Branch, Islamic Azad University, Arak, Iran

ABSTRACT

Pseudomonas aeruginosa is a gram-negative bacterium that is abundant in a variety of moist environments, such as soil and water, and can adapt to many other environments. As an opportunistic pathogen, *Pseudomonas aeruginosa* has been identified as one of the most important hospital pathogens in recent years. It is highly resistant to a variety of antibiotics and antibiotics, and is the main problem of bacterial treatment in important wards of hospitals such as burns and intensive care. In these patients, skin defense is lost. Burn wound healing is a complex process that involves an early burst of energy depletion and cell necrosis, followed by an inflammatory phase, cell death, granulation tissue formation, matrix formation, and collagen rearrangement. Bacterial or antioxidant activity is a good candidate for wound healing. The purpose of this study was to evaluate the effect of thyme essential oil on drug-resistant *Pseudomonas* isolated from cystic fibrosis patients. According to the findings of this study, thyme essential oil has significant antimicrobial activity, and its impact on clinical and drug-resistant *Pseudomonas aeruginosa* can lead to failure of bacterial resistance and accelerate its treatment in clinical wards.

KEYWORDS

Thyme essential oil; Resistance failure; Pseudomonas aeruginosa; Cystic fibrosis.

1. Case Presentation

The Lamiaceae family is one of the largest plant families in the world (excluding the Arctic and Antarctic regions) and has about 200 genera and two to five thousand species of fragrant shrubs and short shrubs. Most mints produce terpenes and a variety of late compounds that store these compounds (mainly) in the epidermal glands of leaves, stems, and reproductive organs (Sheikholeslami, Mousavi, Ashtiani, Doust, & Rezayat, 2016).

Thyme (*Thymus vulgaris* L.) is a genus of dark mint plants that grows as dense shrubs on dry slopes and between boulders in various Mediterranean regions, including France, Portugal, Spain, Italy and Greece. This

plant exists in semi-arid areas of New Zealand for several thousand hectares by car. This species has not been seen wild in our country. Of course, thyme is widely grown every year in Spain, Germany, France, Portugal, the United States, the Czech Republic, Slovakia, Hungary and North Africa. In Iran, the area under cultivation of this species is increasing (Farahani, Barati, Arjomandzadegan, & Vatankhah, 2020).

Thyme is used in a variety of food, pharmaceutical, health and cosmetic industries. Thyme oil has antispasmodic, windbreak, antifungal, disinfectant, anti-worm, anti-dermatitis and expectorant properties. Thyme essential oil is one of the ten famous essential oils that have antibacterial and antifungal properties, antioxidants, natural food preservatives and anti-aging mammals and has a special place in world trade. Thyme is also used in a variety of foods and is used as an aromatic compound in most important food products such as beverages and dairy desserts. Although the growth, quality and quantity of active ingredients in medicinal plants, including dry matter accumulation and essential oil biosynthesis, are primarily controlled by genetic processes, environmental factors also play an important role. What is important is that thyme oil, which is cultivated in different regions, is different in terms of color, taste, viscosity and chemical composition, and seasonal variation has a significant effect on its yield and oil composition (Mottaghiyan, Aghazadeh, Mahmoodzadeh Hosseini, & Imani Fooladi, 2019). Agronomic factors also affect the quantitative and qualitative yield of thyme, which is the planting distance from important agronomic factors that have a significant effect on the yield of essential oil and dry matter per unit area. It was found that shorter plant spacing significantly increased the yield of the branch head and the amount of essential oil, with a planting distance of 5 cm having the highest yield.

Europe, along with the United States, is one of the major consumer markets for thyme. Trade statistics show that the United States imports about 1,000 tons of thyme annually. 90% of thyme oil in world trade is produced in Spain (Farahani et al., 2020).

Pseudomonas bacteria and its characteristics

The common ancestors of the pseudomonad lived hundreds of millions of years ago, yet they were classified by humans in the late nineteenth century. The common name *Pseudomonas* was coined by Walter Migula from 1894 to 1900 as a genus (Chouhan, Sharma, & Guleria, 2017) of gram-negative, rod-shaped, polar, and flagellate bacteria. (*false*) and *monas* means a single unit that can give the meaning of a false unit (*false unit*).

2. Methods

Nutrient agar medium of German Merck company, Blood agar medium of German Merck company and McConkey agar medium of German Merck company and Müllerhinton agar medium of German Merck company were used. The type of bacteria used was *Pseudomonas* and sheep blood was used for blood agar. To perform the tests, the antibiotic discs of chloramphenicol, gentamicin, ampicillin, ciprofloxacin and imipenem, p-paracillin were used. Microliters of thyme essential oil were inoculated on the discs (Basavegowda, Patra, & Baek, 2020; Cunha, 2018; Narasagoudr et al., 2020; RANI, RAO, & SUKUMAR).

The MIC test was performed on a plate of 96 sterile cells using the microdilution broth method (Sardarodiyani, Arianfar, Sani, & Naji-Tabasi, 2019). First, 100 l of Müllerhinton culture medium was poured into 96 microplate wells. Then 100 l of essential oil was added to the first well of each row and diluted from the second to the third house and so on until the ninth house. Finally, 100 l of diluted microbial suspension was added to all wells equivalent to half a McFarland tube. After 24 hours of incubation at 37 ° C by a tray-reading stand base made for this purpose, the bottom of the plate was observed in the mirror under light. We recorded the presence of turbidity, which indicated the growth or non-growth of bacteria, in a special table. According to the concentration definition, the last (thinnest) well was placed in which no turbidity was created. Essential

oil control house, culture medium and microbial equivalent MIC were also considered separately (Sardarodiyani et al., 2019).

3. Results

The results of the study of the susceptibility of Pseudomonas resistant bacteria to thyme essential oil and breaking the resistance of this bacterium are in two stages. In the first stage, the results of the studies performed on the bacterium are presented and the degree of susceptibility is observed. We examine the treatment of rats infected with Pseudomonas.

Initially, the Pseudomonas bacterium isolated from the cystic fibrosis patient, despite clinical tests, was tested for antibiogram by disc diffusion method in the treatment center. The results of determining the antibiotic susceptibility of the isolated bacterium were as follows: the bacterium was evaluated by ciprofloxacin, piperacillin, vimipenem, amikacin, polymyxin B, gentamicin disks, and the obtained growth inhibitions were matched with the CLSI table and the growth inhibition halos were presented as follows (Table 1)

It was observed to be sensitive only to imipenem and semi-sensitive to amikacin and resistant to other antibiotics.

Table 1. Results of Pseudomonas bacterium diffusion disk and conformity of halos to CLSI table

Antibiotic disc	CP	PIP	Polymixin B	AN	GM	IPM
Aura obtained	14	15	5	14	10	24
nterpretation according to CLSI table	S: >21 I:16-20 R: <15	S: >21 I:16-20 R: <14	S: >12 I: -- R: <11	S: >17 I:15-16 R: <14	S: >15 I: 13-14 R: <12	S: >19 I:16-18 R: <15

As explained, to ensure bacterial susceptibility to thyme essential oil, microbial susceptibility tests were performed on isolated Pseudomonas bacteria. These tests include the Kirby Bauer-MIC method, the results of which are presented below.

In Kirby-Bayer method, the amount of growth inhibition halos in different concentrations was investigated and it was observed that in this method up to 0.312 mg / ml growth inhibition halo was observed for Pseudomonas bacteria (Figure 1).



Figure 1. Results of Kirby-Bayer method in bacteria.

Also, the results obtained from the microplate dilution method showed that, a concentration of 5 mg / ml was obtained as a result of MIC (minimum concentration inhibiting bacterial growth).

4. Conclusion

Pseudomonas aeruginosa is a gram-negative bacterium that is abundant in a variety of humid environments such as soil and water and can adapt to many other environments (Alalwan, 2018). This bacterium is also an opportunistic pathogen that infects patients in hospitals, especially burn wards. In these patients, the skin defense is lost. Burn wound healing is a complex process involving an early phase of sudden energy depletion and cellular necrosis that follows the stages of inflammatory phase, cell death, granulation tissue formation, matrix formation, and collagen rearrangement. Drugs that have anti-inflammatory, anti-inflammatory effects Bacterial or antioxidant activity are good candidates for wound healing. The compounds in the extract have all the above effects (Naz, 2018).

From April 1, 2010, Iran and Iran became more and more active due to the presence of uranium and uranium in humans. Expressed. The two had different rates of 0.8% and 0.050%, respectively. From April 1, 2010, Iran and Iran became more and more active due to the presence of uranium and uranium in humans. Expressed. The two had different rates of 0.8% and 0.050%, respectively (Pawar, Bavva, Rohan, & Srivastava, 2020; Sami, 2018).

The results of this study are consistent with previous studies. It is suggested that the extract has antibacterial effects by preventing bacterial growth and multiplication and can also defeat biofilm and its production (Valcourt, 2016).

References

- Alalwan, H. K. A. (2018). Combining nanofabrication with natural antimicrobials to control denture plaque. University of Glasgow.
- Basavegowda, N., Patra, J. K., & Baek, K.-H. (2020). Essential Oils and Mono/bi/tri-Metallic Nanocomposites as Alternative Sources of Antimicrobial Agents to Combat Multidrug-Resistant Pathogenic Microorganisms: An Overview. *Molecules*, 25(5), 1058.
- Chouhan, S., Sharma, K., & Guleria, S. (2017). Antimicrobial activity of some essential oils – present status and future perspectives. *Medicines*, 4(3), 58.
- Cunha, J. A. d. (2018). Efeito antibacteriano do óleo essencial de *Origanum majorana* e de compostos isolados na forma pura e nano encapsulada contra à *Aeromonas hydrophila* em *Rhamdia quelen*. Universidade Federal de Santa Maria.
- Farahani, H., Barati, A., Arjomandzadegan, M., & Vatankhah, E. (2020). Nanofibrous cellulose acetate/gelatin wound dressing endowed with antibacterial and healing efficacy using nanoemulsion of *Zataria multiflora*. *International Journal of Biological Macromolecules*, 162, 762-773.
- Mottaghiyan, Z., Aghazadeh, M., Mahmoodzadeh Hosseini, H., & Imani Fooladi, A. A. (2019). Evaluation of Antibacterial Activity of *Zataria multiflora* Against the Expression of *icaADB* and *aap* Gene and Biofilm Formation in *Staphylococcus epidermidis*. *Archives of Clinical Infectious Diseases*, 14(1).
- Narasagoudr, S. S., Hegde, V. G., Chougale, R. B., Masti, S. P., Vootla, S., & Malabadi, R. B. (2020). Physico-chemical and functional properties of rutin induced chitosan/poly (vinyl alcohol) bioactive films for food packaging applications. *Food Hydrocolloids*, 106096.
- Naz, M. (2018). Synthesis and Characterization of Silver Nanoparticles Using Seed Extracts and their Biomedical Application. Government College University, Lahore.
- Pawar, V., Bavva, M., Rohan, K. V., & Srivastava, R. (2020). Advances in Polysaccharide-Based Antimicrobial Delivery Vehicles Racing for the Surface (pp. 267-295): Springer.
- RANI, G. N., RAO, B. N., & SUKUMAR, E. A PROSPECTIVE STUDY OF HONEY WITH SPECIAL REFERENCE TO ITS ANTIBACTERIAL ACTIVITY.

- Sami, A. (2018). Study on Natural Products and Other Related Compounds Active against Bacteria Associated with Urinary Tract Infections. University of Karachi, Karachi.
- Sardarodiyani, M., Arianfar, A., Sani, A. M., & Naji-Tabasi, S. (2019). Antioxidant and antimicrobial activities of water-soluble polysaccharide isolated from Balangu seed (*Lallemantia royleana*) gum. *Journal of Analytical Science and Technology*, 10(1), 17.
- Sheikholeslami, S., Mousavi, S. E., Ashtiani, H. R. A., Doust, S. R. H., & Rezayat, S. M. (2016). Antibacterial activity of silver nanoparticles and their combination with zataria multiflora essential oil and methanol extract. *Jundishapur journal of microbiology*, 9(10).
- Valcourt, C. (2016). Contribution à l'étude du traitement de bactéries multi-résistantes: Associations de composants aromatiques d'huiles essentiels nano-encapsulés avec des antibiotiques.

Copyrights

Copyright for this article is retained by the author(s), with first publication rights granted to the journal. This is an open-access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/4.0/>).