
Application of Edge Detection Techniques to Model and 3D Field Data

Ofuyah W. N

Department of Earth Sciences, Federal University of Petroleum Resources, Effurun, Nigeria

ABSTRACT

In this study, we present the results of the application of amplitude-derived semblance and cepstral decomposition to model and 3D field data from the Niger Delta. The cepstrum is the Fourier transform of the natural logarithm of the Fourier transform of the data and transforms the data from the frequency to quefrequency domain, while the semblance measures localized trace similarity. Cepstral domain analysis separates source and transmission path effects and provides new improved information on the seal risk of hydrocarbon prospect by revealing the presence of multiple sub-parallel faults capable of increasing the probability of up-dip lateral seal for hydrocarbon traps. This is to guide prospect evaluation in fault – trapped prospect.

KEYWORDS

Fourier transform; Maximum Entropy; Cepstrum; Semblance; Homomorphic; Depobelt.

1. Introduction

Hydrocarbon prospects that are faulted have special risks because faults can cause leakage from hydrocarbon accumulations or provide lateral seal. Data interpretations usually end after identifying the fault plane in the wrong belief that the fault plane itself creates a lateral seal. However an important element of entrapment is the presence of numerous subtle sub-parallel faults. The identification of the faults in the absence of clear break of horizon or diffraction particularly with a computer-based algorithm is a major challenge. (Downey, 1990)

In this study, we present the results of the application of amplitude-derived semblance and cepstral decomposition to model and 3D field data from the Niger Delta. The cepstrum is the Fourier transform of the natural logarithm of the Fourier transform of the data and transforms the data from the frequency to quefrequency domain, while the semblance measures localized trace similarity. Cepstral domain analysis separates source and transmission path effects and provides new improved information on the seal risk of hydrocarbon prospect by revealing the presence of multiple sub-parallel faults capable of increasing the probability of up-dip lateral seal for hydrocarbon traps. This is to guide prospect evaluation in fault –trapped prospect.

1.1 Geologic Setting

The source of our data is the Niger Delta region of which the 'X' -basin is a part is a prolific hydrocarbon province formed during three depositional cycles from middle cretaceous to recent in Nigeria. The recent Niger Delta has the continental, transitional and marine environments comprising braided streams, meander-belt systems, floodplain basins, barrier bars, fire sand, silt and clay. Most of the hydrocarbons are in the sandstones of the Agbada formation, mostly trapped in roll over Anticline, fronting growth faults (Figure 1). The shales create lateral and top seals (Short and Stauble1967). The reservoirs of this formation are typically channels and barrier sandstone bodies and values of porosity and permeability are generally high (up to 40% and 1-2 Darcys respectively).

1.2 Equations

This edge detection attribute is a more meaningful quantity than the coherence attributes (E_t), which measures edges over a narrow window. According to Tanner et al. (1979), semblance (S_t) denotes the ratio of the total energy of the stack, within a gate of length Δt , to the sum of the energy of the component traces within the same time gate. Using the terminology above, we can write semblance

$$S_t = \frac{\sum_{t=\tau}^{t+\Delta t} (\sum_i x_{ti})^2}{\sum_{t=\tau}^{t+\Delta t} \sum_i (x_{ti}^2)} \quad (1)$$

The semblance attribute will not only tend to be large when a coherent event is present but the magnitude of the semblance will also be sensitive to the amplitude of the event. Thus strong events usually exhibit large semblance and weak events will exhibit moderate values of semblance while incoherent data will have very low semblance. (Taner et al., 2009, Ofuyah et al, 2014).

1.3 Cepstral Transform (CT)

Cepstral decomposition extends the widely used process of spectral decomposition. It measures bed thickness even when the bed itself cannot be interpreted (Hall, 2006). Spectral decomposition maps are typically interpreted qualitatively using geomorphologic pattern recognition or semi quantitatively, to infer relative thickness variability Spectral decomposition is rigorous when analyzing subtle stratigraphic plays and fractured reservoirs.

The Cepstrum can be defined as the Fourier transform of the log of the spectrum. Given a noise free trace in time (t) domain as $x(t)$ obtained by convolution of a wavelet and reflectivity series r and assuming $X(f)$, $W(f)$ and $R(f)$ are their frequency domain equivalents, then, Since the Fourier transform is a linear operation, the Cepstrum is

$$F[\ln(\text{mod } X)] = F[\ln(\text{mod } W)] + F[\ln(\text{mod } R)] \quad (2)$$

The Cepstrum is useful because it separates source and filter and can be applied to detect local periodicity. There is a complex cepstrum (Oppenheim, 1965) and a real Cepstrum. The complex Cepstrum method used here is employed to recover signals generated by a convolution process and is called Homomorphic deconvolution (Oppenheim and Schafer, 1968).




Figure 2. Cepstrum and Spectrum compared with original data and log of magnitude of its spectrum (Source: Wikipedia Free Encyclopedia,2017,<https://en.wikipedia.org/wiki/Cepstrum>)

2. Method

The method adopted is fast Fourier transform convolution techniques. The algorithm was tested by computing the semblance attribute of a three-layer geologic model of a fault using a 40Hz Ricker wavelet for simulation. The faulty bed in the model is a low impedance medium embedded in a high impedance medium. The geologic features were analyzed to include stratigraphic and near-vertical structural features by using a long analysis window centered on the top of the fault interval. Cepstral decomposition of the field data was computed by taking the Fourier transform of the natural logarithm of the Fourier transform of the data following the treatment by Hall, 2006.

3. Results and Analysis

The results obtained are shown in Figures 3, 4, 5, and 6. (Figure 3) shows the computed model including its geometry and seismogram. The amplitude and phase spectra, and gamnitude and saphce cepstra are presented as Figure 4, while Figure 5 shows the semblance section using different displays. The semblance attribute will not only tend to be large when a coherent event is present but the magnitude of the semblance will also be sensitive to the amplitude of the event. Thus strong events usually exhibit large semblance and weak events will exhibit moderate values of semblance while incoherent data will have very low semblance. (Taner et al., 2009). In Figure 5, High semblance (Blue) indicates strong and coherent events, weak events exhibit moderate values (Yellow) of semblance, while incoherent data has very low (Green) semblance.

3.1 Application to Model Data




Figure 3(a). Geometry of Geologic Model of a Fault




Figure 3(b). 40 Hz Ricker Wavelet used for the Simulation




Figure 4(a). 40 Hz Fault Model: Spectral




Figure 4(b). 40 Hz Fault Model: Cepstral plots

(b) Amplitude and Phase Spectra of Model. The amplitude recovery here is not discernible.

(c) Gamnitude and Saphe Cepstra of Fault Model. Note the higher amplitude recovery in the cepstral plots and compare with Figure 2. Gamnitude shows discontinuity/sequence boundaries.

Saphe highlights segmentation and lithologic changes.




Figure 5.(a) Fault seismogram displayed with Matlab,

(b) Surfer-8 display of (a)

(c) Semblance of (a)

High semblance (Blue) indicates strong and coherent events. Weak events exhibit moderate values (Yellow) of semblance while incoherent data will have very low (Green) semblance.

3.2 Application to Field Data




Figure 6. Top Reservoir (a) Original amplitude (top) (b) Cepstral gamnitude (bottom)

The Gamnitude shows discontinuity/segmentation and lithologic changes. Four compartments namely, A, B, C and D can be seen in (b) Gamnitude.

4. Conclusions

Faults are critical to the accumulation of hydrocarbon and manifest themselves as abrupt, gradual or gentle changes of seismic amplitude. However an important element of entrapment is the presence of numerous subtle sub-parallel faults and their identification with computer –based algorithm has been a major challenge. In frequency domain, spectral decomposition, requiring the use of a time window to unmask subtle events have high sensitivity to noise, weak frequency resolution arising from applied windows, and are therefore

unreliable. Knowledge of the presence and identification of subtle faults has led to the need for improved techniques. Our main conclusion is that Cepstral decomposition of the thin bed reservoir revealed subtle sub-parallel faults and provided an enhanced level of evaluating the seal risk on prospect. This will facilitate improved reservoir production and performance. The key inputs are the concepts and practices of seismic stratigraphy and principles of spectral and cepstral decomposition, a clear knowledge of signal analysis and properly migrated seismic data.

References

- Doust, H., and Omatsola, E., Niger Delta, in, Edwards, J. D., and Santogrossi, P.A., eds., *Divergent/passive Margin Basins*, American Association of Petroleum Geologists Memoir 48: Tulsa, American Association of Petroleum Geologists, 1990, pp. 239-248.
- Downey, M.W. *Faulting and Hydrocarbon Entrapment*. The Leading Edge; vol. 9; 1990, pp.20-22
- Hall, M. *Predicting Bed Thickness with Cepstral Decomposition*, The Leading Edge, Vol.25, No 2, 2006, pp. 199-204.
- Short, K. C; and Stauble, A. J. *Outline of the Geology of Niger Delta*. American Association of Petroleum Geologists Bulletin, vol. 51, 1967, pp. 761-779.
- Stacher, P., *Present understanding of the Niger Delta hydrocarbon habitat*, in, Oti, M.N., and Postma, G., eds., *Geology of Deltas*: Rotterdam, A.A. Balkema, 1995, pp. 257-267.
- Taner, M.T., Koehler and Sheriff, R. E. *Complex Trace Analysis*, *Geophysics*, (44): 1041-1063. Revisited, 64th Annual International Meeting, SEG, Expanded Abstracts, 1979, 1104-1106.
- Ofuyah, W.N., Alao, O.A., Olaseni, V. and, Adeoti, L. *Seismic Spectral Attributes using Coherence and Semblance Algorithms*, *Computer Engineering and Intelligent Systems*, vol.5, No.9, 2014, pp 11-17
- Oppenheim, A.V. "Superposition in a Class of Nonlinear Systems" Ph.D. diss., Res. Lab. Electronics, MIT, 1965.
- Oppenheim, A. V and Schafer, R. W. "Homomorphic Analysis of Speech", *IEEE Trans. Audio Electro acoust*, vol. AU-16, pp. 221-226, R.W. Schafer, *Echo Removal by Discrete Generalized Linear Filtering*: Res. Lab. Electron. MIT, Tech. Rep., 466, 1969.
- Oppenheim, A. V and Schafer R. W. "Digital Signal Processing", (Prentice Hall), 1975.

Copyrights

Copyright for this article is retained by the author(s), with first publication rights granted to the journal. This is an open-access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/4.0/>).